

Cosmological Magnetic Fields from Primordial Helicity

George B. Field¹ and Sean M. Carroll²

¹ *Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138,*

USA

² *Dept. of Physics and Enrico Fermi Institute, University of Chicago, Chicago, IL 60637,*

USA

Email: gfield@cfa.harvard.edu, carroll@theory.uchicago.edu

Abstract

Primordial magnetic fields may account for all or part of the fields observed in galaxies. We consider the evolution of the magnetic fields created by pseudoscalar effects in the early universe. Such processes can create force-free fields of maximal helicity; we show that for such a field magnetic energy inverse cascades to larger scales than it would have solely by flux freezing and cosmic expansion. For fields generated at the electroweak phase transition, we find that the predicted wavelength today can in principle be as large as ~ 10 kpc, and the field strength can be as large as $\sim 10^{-10}$ G.

The origin of galactic and intergalactic magnetic fields is an unsolved problem [1]. The standard α - Ω dynamo theory of galactic magnetic fields [2] has been criticized for not adequately taking into account the back reaction of the growing magnetic field, and in any case, the theory requires a seed field of unknown origin. There is now an extensive literature examining the possibility that magnetic fields were created by exotic processes in the early universe, although most such processes, if they work at all, produce fields on scales too small to be of interest to astronomy [3].

In magnetohydrodynamics, energy can be transferred from small to large scales by a process known as the inverse cascade. As shown in pioneering work by Pouquet and collaborators [4], a critical ingredient of the inverse cascade mechanism is the presence of substantial magnetic helicity, also known as the Chern-Simons term (although non-helical cascades have also been investigated [5]). The idea that magnetic helicity may drive an inverse cascade from microphysical magnetic fields to large-scale cosmological fields has been advocated by Cornwall [6], and investigated subsequently by Son [7], who proposed scaling properties we will verify below. The present work has two goals: 1.) To work towards an analytic understanding of the cascade process as an initial-value problem appropriate for cosmology, by studying similarity solutions of the MHD equations in the presence of helicity; and 2.) To apply this understanding to proposed mechanisms which create helical primordial fields via pseudoscalar processes in the early universe. A preliminary account has been given in [8].

The early-universe processes we consider can be thought of as arising from the evolution of an electrically neutral pseudoscalar field ϕ , coupled to electromagnetism through a Lagrange density

$$\mathcal{L} = -\frac{1}{4}\phi F_{\mu\nu}\tilde{F}^{\mu\nu} = \phi \mathbf{E} \cdot \mathbf{B} , \quad (1)$$

where $\tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}$ is the dual electromagnetic field strength tensor. (For simplicity we elide the distinction between electromagnetic and hypercharge fields; they are related by factors of order unity.) The pseudoscalar in question may represent an axion [9], a more general pseudo-Goldstone boson [10], or may model the effect of a chemical potential for right-handed electron number (in which case the chemical potential μ is given by the time derivative of ϕ). This last scenario has been proposed by Joyce and Shaposhnikov [11], who showed that it could lead to fields at the electroweak phase transition with magnitude $B_{\text{EW}} \sim 10^{22}$ G and coherent over length scales $\lambda_{\text{EW}} \sim 10^{-8} H_{\text{EW}}^{-1} \sim 2 \times 10^{-9}$ cm.

The field equations for electromagnetism in the presence of the interaction (1) are (in

units with $c = 1$)

$$\begin{aligned}
\partial_t \mathbf{E} &= \nabla \times \mathbf{B} - \mathbf{J} - \dot{\phi} \mathbf{B} - \nabla \phi \times \mathbf{E} \\
\nabla \cdot \mathbf{E} &= \rho_E + \nabla \phi \cdot \mathbf{B} \\
\partial_t \mathbf{B} &= -\nabla \times \mathbf{E} \\
\nabla \cdot \mathbf{B} &= 0 ,
\end{aligned} \tag{2}$$

along with Ohm's law, $\mathbf{J} = \sigma(\mathbf{E} + \mathbf{v} \times \mathbf{B})$, where $\mathbf{v}$ is the velocity of the fluid. In the cosmological context of interest here, ϕ is effectively homogeneous ($\nabla \phi = 0$) and the charge density ρ_E vanishes. The magnetic field then satisfies

$$(\nabla^2 - \partial_t^2) \mathbf{B} = \sigma[\partial_t \mathbf{B} - \nabla \times (\mathbf{v} \times \mathbf{B})] - \dot{\phi} \nabla \times \mathbf{B} . \tag{3}$$

For the case that $\sigma = 0$ it has been shown that solutions of (3) can be unstable to exponential growth in the magnetic field [12]; Garretson, Field and Carroll [10] considered field production by such a mechanism during inflation, concluding that the resulting amplitudes were too small to be of astrophysical interest. If one ignores the displacement current $\partial_t \mathbf{E}$ and sets $\dot{\phi} = 0$, (3) reduces to the induction equation of dissipative MHD. In the situation of interest here the conductivity is non-negligible; the electric field changes slowly on the timescales of interest, so we neglect $\partial_t \mathbf{E}$; and the bulk velocity of the fluid is small, so we neglect $\mathbf{v}$ as well. Hence the appropriate form of (3) is

$$(\partial_t - \eta \nabla^2) \mathbf{B} = \eta \dot{\phi} \nabla \times \mathbf{B} , \tag{4}$$

where $\eta = 1/\sigma$ is the resistivity.

It is useful at this point to go to Fourier space and decompose $\mathbf{B}(\mathbf{k})$ into modes of definite helicity (or equivalently, circular polarization), $\mathbf{B}(\mathbf{k}) = B_+ \hat{u}_+ + B_- \hat{u}_-$; here $\hat{u}_\pm = \hat{u}_1 \pm i \hat{u}_2$, where $\hat{u}_1$, $\hat{u}_2$, and $\hat{u}_3 = \mathbf{k}/k$ form a right-handed orthonormal basis. Then solutions to (4) are of the form

$$B_\pm(\mathbf{k}, t) = B_\pm(\mathbf{k}, 0) \exp \left[-\eta \int k(k \mp \dot{\phi}) dt \right] , \tag{5}$$

For $\dot{\phi}$ positive, the B_+ modes will grow exponentially if k is less than $\dot{\phi}$, with maximum growth rate for $k = \dot{\phi}/2$, while the B_- modes decay away. When $\dot{\phi} = 0$ the field undergoes Ohmic decay, although Joyce and Shaposhnikov show that this effect is unimportant for the relevant wavelengths in their scenario [11].

The expectation value of the magnetic energy density of an isotropic plasma in a volume V can be expressed in terms of an energy spectrum e_k^M as

$$E^M = \frac{1}{2} V^{-1} \int_V \mathbf{B}^2 d^3x = \int_0^\infty e_k^M dk , \quad (6)$$

where

$$e_k^M = 2\pi k^2 \langle \mathbf{B}(\mathbf{k}) \cdot \mathbf{B}^*(\mathbf{k}) \rangle . \quad (7)$$

Similarly, the expectation value of the magnetic helicity (or Chern-Simons) density can be written

$$H^M = V^{-1} \int_V \mathbf{A} \cdot \mathbf{B} d^3x = \int_0^\infty h_k^M dk , \quad (8)$$

where

$$h_k^M = 4\pi k^2 \langle \mathbf{A}(\mathbf{k}) \cdot \mathbf{B}^*(\mathbf{k}) \rangle . \quad (9)$$

Note that while e_k^M is non-negative, h_k^M can be of either sign. The helicity and energy spectra satisfy a realizability condition,

$$|h_k^M| \leq 2k^{-1} e_k^M . \quad (10)$$

We say the field is “maximally helical” if, for every k , h_k^M is of the same sign and saturates this inequality. (See [13] for further discussion.)

In Coulomb gauge ($\mathbf{k} \cdot \mathbf{A} = 0$), the modes of the vector potential and the magnetic field are related by $B_\pm = \pm k A_\pm$. The energy and helicity spectra are then

$$\begin{aligned} e_k^M &= 4\pi k^2 (|B_+|^2 + |B_-|^2) , \\ h_k^M &= 8\pi k (|B_+|^2 - |B_-|^2) . \end{aligned} \quad (11)$$

Since the B_+ modes are amplified and the B_- modes suppressed by the evolution of ϕ , only B_+ will contribute; such a field satisfies $h_k^M = 2k^{-1} e_k^M$, and is therefore maximally helical.

If the spectrum of a maximally helical magnetic field is strongly peaked around some wavenumber k_p , the configuration will be substantially force-free: $\mathbf{J} \times \mathbf{B} = (\nabla \times \mathbf{B}) \times \mathbf{B} = 0$. This can be seen by considering

$$\nabla \times \mathbf{B}(\mathbf{x}) = \int k B_+(\mathbf{k}) \hat{u}_+ e^{i\mathbf{k} \cdot \mathbf{x}} d^3k \sim k_p \mathbf{B}(\mathbf{x}) . \quad (12)$$

Force-freedom has been verified in numerical simulations [14]. This condition plays an important role in the evolution of the fields. The velocities associated with force-free fields are small, protecting them from the Silk damping (photon viscosity) discussed in [15] (which implies in addition that they will not produce detectable distortions in the spectrum of the cosmic microwave background [16]).

We turn now to the principles behind the inverse cascade. Pouquet *et al.* [4] studied MHD turbulence in the eddy-damped quasi-normal Markovian (EDQNM) approximation, in which eddy damping is used to close the nonlinear equations in Fourier space. These equations preserve the ideal invariants of total energy and magnetic helicity. Pouquet *et al.* studied the evolution of the spectra e_k^M , h_k^M , and e_k^V (the kinetic energy) under various conditions. The presence of magnetic helicity causes a shift of magnetic energy from large k to small, as the system attempts to minimize magnetic energy while conserving magnetic helicity [equation (10)]. In their study of greatest interest here, illustrated in Figure 4 of [4], nonhelical kinetic energy, together with maximally helical magnetic energy, are injected at $k = 1$ at a constant rate, and h_k^M is computed for $0.016 < k < 16$ at times $0 < t < 200$. The result is a well-defined wave of magnetic energy and helicity that propagates from $k = 1$ downward, reaching $k = 0.03$ at $t = 200$. Similar results have been obtained elsewhere [14,17,18].

Here we attempt to quantify this phenomenon analytically so as to generalize it to the cosmological case, in which magnetic energy and helicity are injected in a short time. To do so, we use equation (3.4) of [4]:

$$(\partial_t h_k^M)_{\text{Nonlocal}} = k^{-1} \Gamma_k (h_k^V - k^2 h_k^M) + \tilde{\Gamma}_k e_k^M + (\alpha_k^V - \alpha_k^M) e_k^M - 2\nu_k^V k^2 h_k^M. \quad (13)$$

This equation is obtained by expanding the full EDQNM equations at a given k to isolate those nonlinear interactions which are nonlocal in that they involve either $k' < ak$ or $k' > k/a$, where a is a small parameter. The first term in (13), with the transport coefficient in k - space Γ_k , represents the so-called Alfvén effect, in which a relatively strong magnetic field at $k' \ll k$ provides a background on which excitation at k propagates as an Alfvén wave, bringing e_k^M into equipartition with e_k^V , and $k^2 h_k^M$ into equipartition with the kinetic helicity h_k^V . The second term is a correction to the Alfvén effect. The third term represents the α -effect of mean-field dynamo theory, according to which any inequality between the kinetic helicity and $k^2 h_k^M$, the current helicity, at $k' \gg k$, tends to destabilize h_k^M and e_k^M at k . We describe this effect, which is believed to be responsible for stellar magnetic fields

[19], further below. The last term, turbulent viscosity, can be shown to be negligible in our case because the velocities are small.

The third term is of greatest significance here. Particularly important is the appearance of the current helicity $\langle \mathbf{b} \cdot \nabla \times \mathbf{b} \rangle$, whose spectrum is $k^2 h_k^M$. The current helicity was neglected in early discussions of the α -effect, and was first uncovered by Pouquet *et al* [4]. Its importance has been confirmed in [20] by a very different approach, and it plays a crucial role in what follows.

To proceed, we identify the dominant term in (13) by reference to Figures 4 and 6 of [4]. First, note that at the peak k_p of the helicity wave at $t = 200$, we find $e_k^V \approx 0.05 e_k^M$, as would occur if the magnetic field is exerting little force on the fluid. This is consistent with the fact that the injected field is maximally helical, hence substantially force-free according to (12). This view is supported by the fact that in the direct simulation of [14], the configurations formed were still 96% maximally helical (hence substantially force-free) at $t = 45$, with $e_k^V/e_k^M \approx 0.02$ at the peak. It is also consistent with the fact that the field at wavenumbers less than k_p is very weak, and hence unable to bring about equipartition of e_k^V with e_k^M via the Alfvén effect.¹ We conclude that in our case the Alfvén effect is weak, and so we neglect the first term in (13), and with it, the next term, a correction to it. An important implication is that in the helicity wave, the condition of maximal helicity applies:

$$e_k^M = \frac{1}{2} k h_k^M . \quad (14)$$

We are left with the α -term. In light of (14), (13) can be written

$$(\partial_t h_k^M)_{\text{Nonlocal}} = \frac{1}{2} (\alpha_k^V - \alpha_k^M) k h_k^M , \quad (15)$$

demonstrating explicitly the potential of $\alpha_k^V - \alpha_k^M$ to destabilize h_k^M . According to equation (3.6) of [4],

$$\alpha_k^V - \alpha_k^M = -\frac{4}{3} \int_{k/a}^{\infty} \theta_{kqq} (h_q^V - q^2 h_q^M) dq , \quad (16)$$

¹The presence of only a very small velocity in Pouquet's solutions raises the question as to how the field can evolve in time, absent dissipation. The authors are currently investigating this with an approximate model of the Pouquet equation.

where the inverse decay constant θ_{kqq} is defined below [equation (19)]. From Figure 6 of [4], the lack of kinetic energy in the helicity wave means that $|\alpha_k^V| \ll |\alpha_k^M|$, and can be neglected. Thus

$$(\partial_t h_k^M)_{\text{Nonlocal}} = -\frac{1}{2}\alpha_k^M k h_k^M \quad (17)$$

$$= \frac{2}{3}k h_k^M \int_{k/a}^{\infty} \theta_{kqq} q^2 h_q^M dq . \quad (18)$$

Hence, if h_k^M is of one sign throughout the wave (as in our case, where we assume that helicity of only one sign is injected), an instability is possible. Pouquet *et al.* (p. 342) describe how this instability works. Suppose the wave has reached $k_p = k_1$. Because $\alpha_k(k_2)$ is due to helicity at $k > k_2$, and the helicity is stored at k_1 , wavenumbers $k_2 < k_1$ become unstable, so that h_k^M and e_k^M grow there. The growing h_k^M at k_2 destabilizes $k_3 < k_2$, and so forth.

This can only be a crude explanation, as it relies on the application of (13), (15) and (16), which are based solely on nonlocal interactions. The reader should bear in mind, however, that the numerical results in [4], represented in Figures 4, 5, and 6 of that paper, are based on the full EDQNM equations, not just the nonlocal version in (13).

The inverse decay constant θ_{kqq} is given by

$$\theta_{kqq} = (\mu_k + 2\mu_q)^{-1} , \quad (19)$$

where the eddy-damping rate μ_k is given by

$$\begin{aligned} \mu_k = & (\nu + \eta)k^2 + 0.26 \left(\int_0^k p^2 e_p dp \right)^{1/2} \\ & + \sqrt{\frac{2}{3}}k \left(\int_0^k e_p dp \right)^{1/2} . \end{aligned} \quad (20)$$

The first term represents damping by viscosity ν and resistivity η . In astrophysical applications ν and η are typically of order 10^{-6} times the succeeding terms, so this effect is important only at very large wavenumbers; we will ignore it henceforth. The second term parameterizes damping due to self-distortion, and the third that due to the nonlinear interaction of Alfvén waves.

In light of (14), we can use (23) to express (20) as

$$\mu_k = (H^M k_p^3)^{1/2} F(\xi) , \quad (21)$$

where

$$F(\xi) = 0.26 \left(\int_0^\xi \zeta^3 s(\zeta) d\zeta \right)^{1/2} + 0.82 \xi \left(\int_0^\xi \zeta s(\zeta) d\zeta \right)^{1/2} . \quad (22)$$

We now look for similarity solutions of the form

$$h_k^M(t) = g(t)s(\xi) , \quad (23)$$

where the shape function $s(\xi)$ depends on the wavenumber scaled to the peak:

$$\xi = \frac{k}{k_p(t)} . \quad (24)$$

We normalize the shape function by $\int s(\xi) d\xi = 1$, which allows us to express the time dependence of (23) as

$$g(t) = \frac{H^M(t)}{k_p(t)} . \quad (25)$$

Given this *ansatz* and $H^M(t)$, a solution will be fully specified by the shape $s(\xi)$ and the peak wavenumber $k_p(t)$.

Given (22), the evolution equation (18) can be written

$$\frac{\dot{H}^M}{H^M} - \frac{\dot{k}_p}{k_p} \left(1 + \xi \frac{s'}{s} \right) = \left(\frac{1}{2} H^M k_p^3 \right)^{1/2} G(\xi) , \quad (26)$$

where $s' \equiv ds/d\xi$ and

$$G(\xi) = \frac{4}{3} \xi \int_{\xi/a}^\infty \frac{\zeta^2 s(\zeta)}{F(\xi) + 2F(\zeta)} d\zeta . \quad (27)$$

Equation (26) is valid for $k \sim k_p$, in particular for $k = k_p(\xi = 1)$, where $s' = 0$ by definition. Thus, $k_p(t)$ solves the differential equation

$$\frac{d}{dt} \ln \left(\frac{H^M}{k_p} \right) = \left(\frac{1}{2} H^M k_p^3 \right)^{1/2} G(1) , \quad (28)$$

where $G(1)$ is a dimensionless constant of order unity. Pouquet *et al.* [4] considered the case $\dot{H}^M = \text{const}$, so $H^M = \dot{H}^M t$, in which case the solution to (28) is

$$k_p(t) = k_p(t_i) t_i / t , \quad (29)$$

provided that

$$G(1) = \left[\frac{1}{8} \dot{H}^M k_p^3(t_i) t_i^3 \right]^{-1/2} . \quad (30)$$

The numerical solution of the exact EDQNM equations found in [4] fits (29)-(30) very well.

Reassured that we can find similarity solutions consistent with the numerical results, we turn to the $\dot{H}^M = 0$ case, relevant to cosmology once the pseudoscalar ϕ has stopped evolving. Then (26) implies that similarity solutions will once again exist, with

$$k_p(t) = k_p(t_i) \left(\frac{t_i}{t} \right)^{2/3} , \quad (31)$$

provided that

$$G(1) = \left[\frac{9}{8} H^M k_p^3(t_i) t_i^2 \right]^{-1/2} . \quad (32)$$

Note that (31) verifies a result of Son [7], derived from a different line of argument.

In addition to the peak wavelength, we also want to know the rms magnetic field, $B_{\text{rms}} = \langle \mathbf{B}^2 \rangle^{1/2}$. For our maximally helical fields we have

$$\begin{aligned} \langle \mathbf{B}^2 \rangle &= 2 \int_0^\infty e_k^M dk = \int_0^\infty k h_k^M dk \\ &= H^M k_p(t) \int_0^\infty \xi s(\xi) d\xi . \end{aligned} \quad (33)$$

The integral in the last expression is independent of time. Thus, when $H^M = \text{constant}$, (31) implies

$$B_{\text{rms}}(t) = B_{\text{rms}}(t_i) \left(\frac{t_i}{t} \right)^{1/3} . \quad (34)$$

The preceding discussion has assumed a flat spacetime background; it is straightforward to adapt these results to an expanding Robertson-Walker spacetime with metric $ds^2 = -dt^2 + R^2(t)d\mathbf{x}^2 = R^2(t^*)[-dt^{*2} + d\mathbf{x}^2]$, where R is the scale factor and t^* is the conformal time. In the radiation-dominated era, the complete set of MHD equations is conformally invariant (see for example [21]), and our results will apply directly to the conformally transformed quantities

$$B_{\text{rms}}^* = R^2 B_{\text{rms}} , \quad k_p^* = R k_p . \quad (35)$$

Thus, the inverse cascade will be characterized by $k_p^*(t^*) \propto (t^*)^{-2/3}$ and $B_{\text{rms}}^*(t^*) \propto (t^*)^{-1/3}$, where the conformal time in the radiation-dominated era is given by $t^* = 2(t_{\text{EQ}}^{1/2}/R_{\text{EQ}})t^{1/2}$, in which the subscript EQ refers to the epoch of matter-radiation equality. The physical quantities at this epoch are therefore related to their initial values by

$$\begin{aligned}
B_{\text{rms}}(t_{\text{EQ}}) &= R_{\text{EQ}}^{-2} B_{\text{rms}}^*(t_{\text{EQ}}^*) \\
&= R_{\text{EQ}}^{-2} \left(\frac{t_i^*}{t_{\text{EQ}}^*} \right)^{1/3} B_{\text{rms}}^*(t_i^*) \\
&= \frac{R^2(t_i)}{R_{\text{EQ}}^2} \left(\frac{t_i}{t_{\text{EQ}}} \right)^{1/6} B_{\text{rms}}(t_i)
\end{aligned} \tag{36}$$

and

$$\begin{aligned}
k_p(t_{\text{EQ}}) &= R_{\text{EQ}}^{-1} k_p^*(t_{\text{EQ}}^*) \\
&= R_{\text{EQ}}^{-1} \left(\frac{t_i^*}{t_{\text{EQ}}^*} \right)^{2/3} k_p^*(t_i^*) \\
&= \frac{R(t_i)}{R_{\text{EQ}}} \left(\frac{t_i}{t_{\text{EQ}}} \right)^{1/3} k_p(t_i) .
\end{aligned} \tag{37}$$

It is possible that additional inverse cascade occurs in the matter-dominated era following t_{EQ} . Pending further study, we will assume here that the effect of this is negligible, and so the following estimates of the characteristic scale of the field today are lower limits. Thus, for $t > t_{\text{EQ}}$ the field is frozen in, the wavenumber scales as R^{-1} , and the magnetic field as R^{-2} . Their values today are thus

$$B_{\text{rms}}(t_0) = \frac{R^2(t_i)}{R_0^2} \left(\frac{t_i}{t_{\text{EQ}}} \right)^{1/6} B_{\text{rms}}(t_i) \tag{38}$$

and

$$k_p(t_0) = \frac{R(t_i)}{R_0} \left(\frac{t_i}{t_{\text{EQ}}} \right)^{1/3} k_p(t_i) . \tag{39}$$

In summary, a maximally helical primordial magnetic field created at time t_i (during radiation domination), with initial amplitude $B_{\text{rms}}(t_i)$ and initial coherence length $\lambda(t_i) = 2\pi/k_p(t_i)$, will undergo an inverse cascade, increasing its length scale by a factor $(t_{\text{EQ}}/t_i)^{1/3}$ over and above stretching due to the expansion of the universe, while its amplitude is diluted by an additional factor $(t_i/t_{\text{EQ}})^{1/6}$. Note that our estimates of the present coherence length substantially exceed those of Son [7], who terminated the inverse cascade at e^+/e^- annihilation because of increasing viscosity. We have argued that our modes are nearly force-free, with negligible velocities; viscosity has no effect.

For purposes of illustration, let us consider the fate of a magnetic field created at the electroweak phase transition ($T_{\text{EW}} = 200$ GeV), so $t_i = t_{\text{EW}} = 6 \times 10^{-12}$ sec. We express the initial coherence length of the field in terms of the Hubble radius, $\lambda(t_{\text{EW}}) = f_\lambda H_{\text{EW}}^{-1} = 0.4 f_\lambda$ cm, and the initial amplitude in terms of the total energy density, $B_{\text{rms}}(t_{\text{EW}}) = f_B \sqrt{8\pi\rho_{\text{EW}}} =$

$2 \times 10^{25} f_B$ Gauss, where f_λ and f_B are dimensionless factors less than unity. (We have switched here from Lorentz-Heaviside units to CGS in order to express the magnetic field in Gauss.) The field today will be coherent over scales

$$\lambda(t_0) = 5 \times 10^{22} f_\lambda \text{ cm} = 20 f_\lambda \text{ kpc} , \quad (40)$$

with amplitude

$$B_{\text{rms}}(t_0) = 4 \times 10^{-10} f_B \text{ Gauss} . \quad (41)$$

In the electroweak case, then, the characteristic length scale of the field has been amplified by a factor of $(t_{\text{EQ}}/t_{\text{EW}})^{1/3} = 6 \times 10^7$ more than would be expected for a frozen-in configuration. If the initial field is coherent over the Hubble radius at the electroweak scale ($f_\lambda \sim 1$) and comparable in energy to the total energy ($f_B \sim 1$), this results in a length scale of ~ 20 kpc and an amplitude of $\sim 10^{-9}$ Gauss. According to Dolag *et al.* [22], the primordial field required to explain Faraday rotation measures of the Coma cluster is $\sim 10^{-9}$ Gauss. The observations are consistent with scales of ~ 60 kpc. Thus, if $f_\lambda \sim f_B \sim 1$, primordial helicity could explain the fields in the Coma cluster.

However, Joyce and Shaposhnikov [11] have estimated that their scenario for helical field generation at the electroweak scale results in fields with $f_\lambda \sim 10^{-8}$ and $f_B \sim 2 \times 10^{-3}$. While intriguing (for example as a candidate seed field for a galactic dynamo), these parameters fall short of providing a sufficient explanation for the fields seen in galaxies today. It is therefore worth considering variations on this mechanism, perhaps with different dynamics for the pseudoscalar ϕ , or field creation at a later epoch such as the QCD scale. Mechanisms which create large-amplitude fields without appreciable helicity will undergo significantly less (if any) inverse cascade, and thus have a difficult time leading to fields of astrophysical significance in the present universe.

ACKNOWLEDGMENTS

This work was supported in part by the National Science Foundation under grant PHY/94-07195 and NASA under grant NAGW-931, and GF acknowledges discussions at the Institute for Theoretical Physics Program on Astrophysical Turbulence, April-June, 2000.

REFERENCES

- [1] For reviews see E.G. Zweibel and C. Heiles, *Nature* **385**, 131 (1997); A.V. Olinto, in *Proceedings of the 3rd RESCEU Symposium*, University of Tokyo, Nov. 10-13, 1997, Ed. K. Sato, T. Yanagida, and T. Shiromizu; [astro-ph/9807051](#).
- [2] A.A. Ruzmaikin, A.M. Shukurov, and D.D. Sokoloff, *Magnetic Fields of Galaxies*, Kluwer, Dordrecht (1988).
- [3] O. Törnqvist, [astro-ph/0002307](#).
- [4] A. Pouquet, U. Frisch, and J. Léorat, *J. Fluid Mech.* **77**, 321 (1976).
- [5] A. Brandenburg, K. Enqvist and P. Olesen, *Phys. Rev. D* **54**, 1291 (1996), [hep-ph/9608422](#); P. Olesen, [astro-ph/9610154](#); K. Enqvist, [astro-ph/9707300](#); T. Shiromizu, [astro-ph/9810339](#).
- [6] J.M. Cornwall, *Phys. Rev. D* **56**, 6146 (1997); [hep-th/9704022](#).
- [7] D.T. Son, *Phys. Rev. D*, **59**, 063008 (1999); [hep-ph/9803412](#).
- [8] S.M. Carroll and G.B. Field, in *33rd Rencontres de Moriond: Fundamental Parameters in Cosmology*, 17-24 January 1998, Les Arcs, France; [astro-ph/9807159](#).
- [9] M.S. Turner and L.M. Widrow, *Phys. Rev. D* **37**, 2743 (1988).
- [10] W.D. Garretson, G.B. Field, and S.M. Carroll, *Phys. Rev. D* **46**, 5346 (1992); [hep-ph/9209238](#).
- [11] M. Joyce and M. Shaposhnikov, *Phys. Rev. Lett.* **79**, 1193 (1997); [astro-ph/9703005](#).
- [12] S.M. Carroll, G.B. Field, and R. Jackiw, *Phys. Rev. D* **41**, 1231 (1990); A.A. Andrianov, R. Soldati, and L. Sorbo, *Phys. Rev. D* **59**, 025002 (1999); [hep-th/9806220](#).
- [13] U. Frisch, A. Pouquet, J. Léorat, and A. Mazure, *J. Fluid Mech.* **68**, 769 (1975).
- [14] M. Meneguzzi, U. Frisch, and A. Pouquet, *Phys. Rev. Lett.* **47**, 1060 (1981).
- [15] K. Jedamzik, V. Katalinić, and A.V. Olinto, *Phys. Rev. D* **57**, 3264 (1998); [astro-ph/9606080](#).
- [16] K. Jedamzik, V. Katalinić, and A.V. Olinto, *Phys. Rev. Lett.* **85**, 700 (2000); [astro-ph/9911100](#).

- [17] D. Balasara and A. Pouquet, *Phys. Plasmas* **6**, 89 (1999).
- [18] A. Brandenburg, [astro-ph/0006186](#).
- [19] H.K. Moffatt, *Magnetic Field Generation in Electrically Conducting Fluids* (Cambridge U. Press, Cambridge), 1978.
- [20] G.B. Field, E.G. Blackman, and H. Chou, *Astrophys. Journ.* **513**, 638 (1999); [astro-ph/9810345](#).
- [21] K. Subramanian and J.D. Barrow, *Phys. Rev. D* **58**, 083502 (1998); [astro-ph/9712083](#).
- [22] K. Dolag, M. Bartelmann, and H. Lesch, [astro-ph/9906329](#).